

Coherent source radius in $p\bar{p}$ collisions

Q. H. Zhang

Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany

X. Q. Li

China Center of Advanced Science and Technology (World Laboratory), P.O. Box 8730, Beijing

100080, P.R. China

Physics Department, Nankai University, Tianjing 300071, P.R. China

(February 1, 2008)

Abstract

We use a recently derived result to extract from two-pion interferometry data from $p\bar{p}$ collisions the radius of the coherent component in the source.

We find a coherent source radius of about $2fm$.

PACS numbers: 25.75. -q, 25.70. Pq, 25.70 Gh.

Two-particle Bose-Einstein interferometry (also known as Hanbury Brown-Twiss intensity interferometry) as a method for obtaining information on the space-time geometry and dynamics of high energy collisions has recently received intensive theoretical and experimental attention. Detailed investigations have shown that high-quality two-particle correlation data can reveal not only the geometric extension of the particle-emitting source but also its dynamical state at particle freeze-out [1–6].

For a partially coherent source, it was suggested that two-pion interferometry can give the coherent degree information of the source [7–9]. In paper [10], we have generalized the two-pion interferometry formula for a partially coherent source and found that two-pion interferometry was also sensitive to the size of the coherent source component. The

physical reason is clear: The particle spectrum distribution is the Fourier transformation of the source distribution, the coherent source radius information must hide in the spectrum distribution and also must appear in the two-pion interferometry formula. We will shown below that a general two-pion interferometry formula satisfy the needs to reveal both the degree of coherence and space-time information of the coherent source which was neglected in previous publications [9,11–14].

For a partially coherent source, one can decompose the pion current into a coherent and a chaotic term [8,10,16]:

$$J(x) = J_c(x) + J_I(x), \quad (1)$$

with

$$J_I(x) = \int d^4x' d^4p j(x', p) \gamma(x') \exp(ip(x - x')). \quad (2)$$

Here $j(x', p)$ is the probability amplitude of finding a pion with momentum p , emitted by the emitter at x' . $\gamma(x') = \exp(i\phi(x'))$ is a random phase factor which has been taken away from $j(x', p)$. All emitters are uncorrelated when assuming:

$$\langle \gamma^*(x) \gamma(y) \rangle = \delta^4(x - y). \quad (3)$$

Here $\langle \dots \rangle$ means phase average. Then the probability amplitude in momentum space reads:

$$\begin{aligned} \tilde{J}(p) &= \int J(x) \exp(-ip \cdot x) d^4x \\ &= \tilde{J}_I(p) + \tilde{J}_c(p), \end{aligned} \quad (4)$$

with

$$\tilde{J}_I(p) = \int J_I(x) \exp(-ip \cdot x) d^4x, \quad \tilde{J}_c(p) = \int J_c(x) \exp(-ip \cdot x) d^4x. \quad (5)$$

Using the relationship

$$\langle J_I^*(x) J_c(y) \rangle = 0, \quad (6)$$

we have

$$< \tilde{J}_I^*(p) \tilde{J}_c(p) > = 0. \quad (7)$$

Then the single pion spectrum can be expressed as:

$$\begin{aligned} P_1(p) &= < \tilde{J}^*(p) \tilde{J}(p) > \\ &= < \tilde{J}_I^*(p) \tilde{J}_I(p) > + \tilde{J}_c^*(p) \tilde{J}_c(p). \end{aligned} \quad (8)$$

Using relationship

$$\begin{aligned} < \gamma^*(x_1) \gamma^*(y_1) \gamma(x_2) \gamma(y_2) > &= < \gamma^*(x_1) \gamma(x_2) > < \gamma^*(y_1) \gamma(y_2) > \\ &+ < \gamma^*(x_1) \gamma(y_2) > < \gamma^*(y_1) \gamma(x_2) >, \end{aligned} \quad (9)$$

we obtain

$$< \tilde{J}_I^*(p_1) \tilde{J}_I^*(p_2) \tilde{J}_I(p_2) \tilde{J}_I(p_1) > = < |\tilde{J}_I(p_1)|^2 > \cdot < |\tilde{J}_I(p_2)|^2 > + |< \tilde{J}_I^*(p_1) \tilde{J}_I(p_2) >|^2. \quad (10)$$

Performing the phase average and using Eq.(4), Eq.(8) and Eq.(10), we can write the two-pion spectrum as

$$\begin{aligned} P_2(p_1, p_2) &= < \tilde{J}^*(p_1) \tilde{J}^*(p_2) \tilde{J}(p_2) \tilde{J}(p_1) > \\ &= P_1(p_1) P_1(p_2) + |< \tilde{J}_I^*(p_1) \tilde{J}_I(p_2) >|^2 \\ &+ 2Re(< \tilde{J}_I^*(p_1) \tilde{J}_I(p_2) > \tilde{J}_c^*(p_2) \tilde{J}_c(p_1)). \end{aligned} \quad (11)$$

Similar to ref. [10], we define the following functions:

$$g_I(x, k) = \int d^4y < J_I^*(x + y/2) J_I(x - y/2) > \exp(ik \cdot y), \quad (12)$$

$$g_c(x, k) = \int d^4y J_c^*(x + y/2) J_c(x - y/2) \exp(ik \cdot y). \quad (13)$$

Here $g_I(x, p)$ and $g_c(x, p)$ are Wigner functions which describes the source distribution of the chaotic source and the coherent source respectively [10,15,16]. From the above definitions, we have:

$$\rho_I(i, j) = \rho_I(p_i, p_j) = \langle \tilde{J}_I^*(p_i) \tilde{J}_I(p_j) \rangle = \int g_I(x, \frac{p_i + p_j}{2}) \exp(i(p_i - p_j) \cdot x) d^4x \quad (14)$$

and

$$\rho_c(i, j) = \rho_c(p_i, p_j) = \tilde{J}_c^*(p_i) \tilde{J}_c(p_j) = \int g_c(x, \frac{p_i + p_j}{2}) \exp(i(p_i - p_j) \cdot x) d^4x. \quad (15)$$

Then the two-pion correlation function for a partially coherent source can be expressed as:

$$C_2(p_1, p_2) = \frac{P_2(p_1, p_2)}{P_1(p_1) \cdot P_1(p_2)}, \quad (16)$$

with

$$P_2(p_1, p_2) = P_1(p_1) \cdot P_1(p_2) + |\rho_I(1, 2)|^2 + 2\text{Re}(\rho_I(1, 2)\rho_c(2, 1)) \quad (17)$$

and

$$P_1(p) = \rho_I(p, p) + \rho_c(p, p). \quad (18)$$

For simplicity [8], we assume that $\rho_I(i, j)$ and $\tilde{J}_c(p)$ are real, then Eq. (16) can be re-written as [9,8]

$$C_2(p_1, p_2) = 1 + \epsilon(p_1)\epsilon(p_2)H_I^2(p_1, p_2) + 2\sqrt{\epsilon(p_1)\epsilon(p_2)(1 - \epsilon(p_1))(1 - \epsilon(p_2))}H_I(p_1, p_2) \quad (19)$$

with

$$H_I(p_1, p_2) = \frac{\rho_I(1, 2)}{\sqrt{\rho_I(p_1, p_1)\rho_I(p_2, p_2)}}, \epsilon(p) = \frac{\rho_I(p, p)}{P_1(p)}. \quad (20)$$

Here $\epsilon(p)$ describes the chaotic degree of the source. For a totally chaotic source, $\epsilon(p) = 1$ while for a totally coherent source, $\epsilon(p) = 0$. Due to the limited statistics, the dependence of $\epsilon(p)$ on momentum p is often neglected, then the above formula can be simplified as [9,11,12]

$$C_2(p_1, p_2) = 1 + \epsilon^2 H_I^2(p_1, p_2) + 2\epsilon(1 - \epsilon)H_I(p_1, p_2). \quad (21)$$

Eq. (21) is now widely used in the literature. In the following, we will show that Eq. (21) is too simple to approximate Eq.(19) and the information about the radius of the

coherent source is lost in Eq.(21). We will derive a formula which is better than Eq.(21) to approximate Eq. (19). Eq. (16) can be re-written as

$$C_2(p_1, p_2) = 1 + \frac{|\rho_I(1, 2)|^2}{P_1^2(k)} \cdot \frac{P_1^2(k)}{P_1(p_1)P_1(p_2)} + 2\text{Re} \left(\frac{\rho_c(1, 2)\rho_I(2, 1)}{P_1^2(k)} \right) \frac{P_1^2(k)}{P_1(p_1)P_1(p_2)}. \quad (22)$$

Here $k = (p_1 + p_2)/2$. Because the single particle distribution is exponential form in $p\bar{p}$ and ee^+ collisions [4], we have $P_1^2(k)/P(p_1)/P(p_2) \sim 1$. Under the assumption that $\rho_I(i, j)$ and $\tilde{j}_c(p)$ are real, the above formula can then be re-expressed as

$$C(p_1, p_2) = 1 + \epsilon^2(k)R_I^2(p_1, p_2) + 2\epsilon(k)(1 - \epsilon(k))R_c(2, 1)R_I(1, 2) \quad (23)$$

with

$$R_I(p_1, p_2) = \frac{\rho_I(1, 2)}{\rho_I(k, k)}, R_c(p_1, p_2) = \frac{\rho_c(1, 2)}{\rho_c(k, k)}. \quad (24)$$

One clearly sees the difference between Eq.(23) and Eq.(21), a extra term $R_c(p_1, p_2)$ appears in Eq.(23). Similar to ref. [14], we take both $R_I(p_1, p_2)$ and $R_c(p_1, p_2)$ as Gaussian functions and neglect the dependence of ϵ on momentum, then Eq.(23) can be written down as

$$C_2(p_1, p_2) = 1 + \epsilon^2 \exp(-r_I^2 \cdot q^2) + 2\epsilon(1 - \epsilon) \exp(-(r_I^2 + r_c^2) \cdot q^2/2). \quad (25)$$

Here r_I and r_c is the radius of the chaotic source and the coherent source respectively. Eq.(25) is a limiting form of the general expressions derived in Ref [10] for the case of a 1-dimensional Gaussian parametrization. In the above derivation, we did not take into account the influence of resonance decay on the two-pion correlations which may distort the simple Gaussian function assumed above and may also result in an effective double Gaussian two-pion correlation function. In paper [14], Kulka et al. used the following formula

$$C_2(p_1, p_2) = 1 + \lambda_1 \cdot \exp(-R_1^2 \cdot q^2) + \lambda_2 \exp(-R_2^2 \cdot q^2) \quad (26)$$

to fit their data and found that $\lambda_1 = 0.30_{-0.09}^{+0.08}$, $\lambda_2 = 0.45 \pm 0.13$, $R_1 = 0.63 \pm 0.04 fm$ and $R_2 = 2.11 \pm 0.63 fm$ with $\chi^2/NDF = 1.09$. In the paper [14], Kulka et al found that Eq.(26) was not consistent with Eq. (21). But if we take those values into Eq.(25), we will find that

$r_I = 0.63 \pm 0.04 \text{ fm}$, $r_c = 2.01 \pm 0.67 \text{ fm}$ and $\epsilon = 0.54 \pm 0.07$. It is very interesting to notice that the size of the coherent source produced in $p\bar{p}$ collisions is around 2 fm which is larger than the radius of chaotic source. Also from the above comparison we found that Eq.(21) is too simple to approximate Eq.(19).

Conclusions: In this paper, it was shown that the widely used two pion interferometry formula (Eq.(21)) was too simple to approximate Eq.(19). The information about the radius of coherent source was lost in previous publication. Using Eq.(23) to fit the data, we found that the radius of coherent source produced in $p\bar{p}$ collisions was around 2 fm .

ACKNOWLEDGMENTS

One of the author (Q.H.Z.) express his gratitude to Drs. U. Heinz, U. Widemann and Y. Pang for helpful discussion. This work was supported by the Alexander von Humboldt Foundation and China national natural Science Foundation.

REFERENCES

- [1] D.H. Boal, C.K. Gelke and B.K. Jennings, Rev. Mod. Phys. V62, 553 (1990).
- [2] I.V. Andreev, M. Pümer and R.M. Weiner, Int. J. Mod. Phys. A8, 4577 (1993).
- [3] S. Pratt, in Quark-Gluon-Plasma 2, ed. R. C. Hwa (World Scientific Publ. Co. Singapore 1995) p. 700.
- [4] U. Heinz; in: *Correlations and Clustering Phenomena in Subatomic Physics*, ed. by M.N. Harakeh, O. Scholten, and J.H. Koch, NATO ASI Series B**359** (Plenum, New York, 1997) p.137.
- [5] W.A. Zajc, in: Hadronic Matter in Collision, edited by P. Carruthers and D. Strottman (World Scientific, Singapore, 1986), p. 43.
- [6] B. Lörstand, Int. J. Mod. Phys. A4, 861 (1989).
- [7] G.N. Fowler and R.M. Weiner, Phys. Rev. D**17**, 3118 (1978).
- [8] M. Gyulassy, S.K. Kauffmann, L. W. Wilson, Phys. Rev. C**20**, 2267 (1979).
- [9] R. M. Weiner, Phys. Lett. B**232**, 278 (1989).
- [10] U. Heinz and Q. H. Zhang, Phys. Rev. C**56**, 426 (1997).
- [11] T. Alexopoulos et. al Phys. Rev. D**48**, 1931 (1993).
- [12] Mark II Collab., I. Juricic et al., Phys. Rev. D**39**, 1(1989).
- [13] UA1-Minimum Bias-Collaboration, N. Neumeister et al, Phys. Lett. B **275**, 186 (1992).
T. Åkesson et al. AFS Coll.: Z. Phys. C **36**, 517 (1987).
- [14] K. Kulka and B. Lörstand, Z. Phys. C **45**, 581 (1990).
- [15] S. Pratt, Phys. Rev. Lett.**53**, 1219 (1984)
- [16] W.Q. Chao, C.S. Gao and Q.H. Zhang; J. Phys. G**21**, 847 (1995).